



# SOUTHEAST GAP ANALYSIS PROJECT



## Species Modeling Report


### Yellow-throated Vireo

*Vireo flavifrons*


Taxa: Avian  
 Order: Passeriformes  
 Family: Vireonidae

SE-GAP Spp Code: **bYTVI**  
 ITIS Species Code: 179009  
 NatureServe Element Code: ABPBW01170

#### KNOWN RANGE:



#### PREDICTED HABITAT:



Range Map Link: [http://www.basic.ncsu.edu/segap/datazip/maps/SE\\_Range\\_bYTVI.pdf](http://www.basic.ncsu.edu/segap/datazip/maps/SE_Range_bYTVI.pdf)

Predicted Habitat Map Link: [http://www.basic.ncsu.edu/segap/datazip/maps/SE\\_Dist\\_bYTVI.pdf](http://www.basic.ncsu.edu/segap/datazip/maps/SE_Dist_bYTVI.pdf)

GAP Online Tool Link: <http://www.gapservice.ncsu.edu/segap/segap/index2.php?species=bYTVI>

Data Download: [http://www.basic.ncsu.edu/segap/datazip/region/vert/bYTVI\\_se00.zip](http://www.basic.ncsu.edu/segap/datazip/region/vert/bYTVI_se00.zip)

#### PROTECTION STATUS:

Reported on March 14, 2011

Federal Status: ---

State Status: ID (P), KY (N), NJ (S/S), NV (YES), NY (PB), RI (Not Listed), UT (None), QC (Non suivie)

NS Global Rank: G5

NS State Rank: AL (S5B), AL (S5B), AR (S4B,S4N), AZ (SNA), CA (SNA), CO (SNA), CT (S5B), CT (S5B), DC (S2S3B,S2S3N), DE (S3B), FL (SNRB), GA (S4), IA (S4B,S4N), ID (SNA), IL (S4S5), IN (S4B), KS (S3B), KY (S5B), LA (S4B), MA (S4B), MD (S4S5B), ME (S3B), MI (S5), MN (SNRB), MO (SNRB), MS (S5B), MS (S5B), MT (SNA), NC (S4B), NC (S4B), ND (SNRB), NE (S3), NH (S4B), NJ (S4B), NM (SNA), NV (SNA), NY (S5), OH (S5), OK (S2B), PA (S4B), RI (S4B), SC (S3?B), SD (S2B), SD (S2B), TN (S4), TX (S4), UT (SNA), VA (S4), VT (S5B), VT (S5B), WA (SNA), WI (S4B), WI (S4B), WV (S5B), WY (SNA), MB (S4B), MB (S4B), NB (SNA), NF (SNA), NS (SNA), ON (S4B), PE (SNA), QC (S4B), SK (S2B,S3N)

**SUMMARY OF PREDICTED HABITAT BY MANAGMENT AND GAP PROTECTION STATUS:**

	US FWS		US Forest Service		Tenn. Valley Author.		US DOD/ACOE	
	ha	%	ha	%	ha	%	ha	%
Status 1	46,510.3	< 1	10,936.2	< 1	0.0	0	0.0	0
Status 2	76,721.6	< 1	114,047.9	< 1	0.0	0	3,593.1	< 1
Status 3	2,003.5	< 1	772,013.4	2	47,847.9	< 1	257,314.1	< 1
Status 4	60.9	< 1	0.0	0	0.0	0	87.4	< 1
Total	125,296.3	< 1	896,997.5	3	47,847.9	< 1	260,994.6	< 1
	US Dept. of Energy		US Nat. Park Service		NOAA		Other Federal Lands	
	ha	%	ha	%	ha	%	ha	%
Status 1	0.0	0	52,887.0	< 1	7.7	< 1	1,686.6	< 1
Status 2	0.0	0	5,645.5	< 1	2,757.4	< 1	13.4	< 1
Status 3	22,212.4	< 1	64,773.4	< 1	0.0	0	2,094.3	< 1
Status 4	0.0	0	0.0	0	0.0	0	0.0	0
Total	22,212.4	< 1	123,305.9	< 1	2,765.1	< 1	3,794.3	< 1
	Native Am. Reserv.		State Park/Hist. Park		State WMA/Gameland		State Forest	
	ha	%	ha	%	ha	%	ha	%
Status 1	0.0	0	790.9	< 1	52.5	< 1	0.0	0
Status 2	0.0	0	5,388.7	< 1	287,529.4	< 1	546.0	< 1
Status 3	13,102.4	< 1	157,134.3	< 1	112,634.7	< 1	67,422.3	< 1
Status 4	0.0	0	0.0	0	43,921.4	< 1	12.1	< 1
Total	13,102.4	< 1	163,313.9	< 1	444,138.0	1	67,980.4	< 1
	State Coastal Reserve		ST Nat.Area/Preserve		Other State Lands		Private Cons. Easemt.	
	ha	%	ha	%	ha	%	ha	%
Status 1	0.0	0	6,310.9	< 1	0.0	0	0.0	0
Status 2	5,223.1	< 1	36,030.2	< 1	5.0	< 1	841.6	< 1
Status 3	0.0	0	4,658.1	< 1	8,044.5	< 1	27,245.9	< 1
Status 4	0.0	0	0.3	< 1	1,837.4	< 1	0.0	0
Total	5,223.1	< 1	46,999.5	< 1	9,886.9	< 1	28,087.5	< 1
	Private Land - No Res.		Water		Overall Total			
	ha	%	ha	%	ha	%		
Status 1	0.0	0	0.0	0	119,182.0	< 1		
Status 2	0.0	0	0.0	0	538,342.9	2		
Status 3	226.4	< 1	0.0	0	1,558,727.6	7		
Status 4	32,608,766.3	91	35,995.6	< 1	32,734,541.8	92		
Total	32,608,992.6	91	35,995.6	< 1	34,950,794.2	100		

GAP Status 1: An area having permanent protection from conversion of natural land cover and a mandated management plan in operation to maintain a natural state within which disturbance events (of natural type, frequency, and intensity) are allowed to proceed without interference or are mimicked through management.

GAP Status 2: An area having permanent protection from conversion of natural land cover and a mandated management plan in operation to maintain a primarily natural state, but which may receive use or management practices that degrade the quality of existing natural communities.

GAP Status 3: An area having permanent protection from conversion of natural land cover for the majority of the area, but subject to extractive uses of either a broad, low-intensity type or localized intense type. It also confers protection to federally listed endangered and threatened species throughout the area.

GAP Status 4: Lack of irrevocable easement or mandate to prevent conversion of natural habitat types to anthropogenic habitat types. Allows for intensive use throughout the tract. Also includes those tracts for which the existence of such restrictions or sufficient information to establish a higher status is unknown.

## PREDICTED HABITAT MODEL(S):

### Summer Model:

Habitat Description: Yellow-throated vireos breed in a variety of edge habitats in bottomland and upland mature deciduous and mixed deciduous forests (Rodewald and James 1996) with little undergrowth (Kaufman 1996). Suitable habitat includes forest edges of streams, rivers, swamps, roads, and open wooded habitats with tall deciduous trees, such as those found in parks, orchards, or small towns. They prefer mature forests such as oak-hickory, oak-gum-cypress, oak-pine, elm-ash-cottonwood, maple-beech-birch, maple-oak-basswood and maple basswood. They generally are not found in pure coniferous stands and do not breed in forest interiors unless there is openings (Rodewald and James 1996). Nests from mid-April to late July (peak mid-May to late June) in the mid-Atlantic region (Bushman and Therres 1988). Clutch size 3-5 (usually 4). Incubation 14-15 days, by both sexes. Young tended by both parents, leave nest at about 15 days. Fledglings are independent in about four weeks, but remain with parents until August (Terres 1980, Harrison 1975, James 1973, Bent 1950). Males establish territories before mating occurs (Nest information from NatureServe 2005). Amy Silvano 21apr05

Ecosystem Classifiers: Mixed, Hardwoods, Mesic, Mesic Cove & Montane Forests, Glades (Woodlands only), Low Urban, Deciduous Plantation, Swamps, Floodplain/Riparian (excluding herb. Modifiers). Amy Silvano 21apr05

Elevation Mask: < 1000m

Mask of Forest/Open Ecotone: Include within 250m of ecotone edge.

### Selected Map Units:

Functional Group	Map Unit Name
Anthropogenic	Deciduous Plantations
Anthropogenic	Developed Open Space
Anthropogenic	Low Intensity Developed
Forest/Woodland	Alabama Ketona Glade and Woodland
Forest/Woodland	Allegheny-Cumberland Dry Oak Forest and Woodland
Forest/Woodland	Allegheny-Cumberland Dry Oak Forest and Woodland - Hardwood Modifier
Forest/Woodland	Appalachian Hemlock-Hardwood Forest
Forest/Woodland	Atlantic Coastal Plain Dry and Dry-Mesic Oak Forest
Forest/Woodland	Atlantic Coastal Plain Fall-line Sandhills Longleaf Pine Woodland - Offsite Hardwood Modifier
Forest/Woodland	Atlantic Coastal Plain Mesic Hardwood and Mixed Forest
Forest/Woodland	Atlantic Coastal Plain Northern Mixed Oak-Heath Forest
Forest/Woodland	Central and Southern Appalachian Montane Oak Forest
Forest/Woodland	Central and Southern Appalachian Northern Hardwood Forest
Forest/Woodland	Central Appalachian Alkaline Glade and Woodland
Forest/Woodland	Central Appalachian Oak and Pine Forest
Forest/Woodland	East Gulf Coastal Plain Interior Shortleaf Pine-Oak Forest - Hardwood Modifier
Forest/Woodland	East Gulf Coastal Plain Interior Shortleaf Pine-Oak Forest - Mixed Modifier
Forest/Woodland	East Gulf Coastal Plain Interior Upland Longleaf Pine Woodland - Offsite Hardwood Modifier
Forest/Woodland	East Gulf Coastal Plain Limestone Forest
Forest/Woodland	East Gulf Coastal Plain Northern Dry Upland Hardwood Forest
Forest/Woodland	East Gulf Coastal Plain Northern Loess Bluff Forest
Forest/Woodland	East Gulf Coastal Plain Northern Loess Plain Oak-Hickory Upland - Hardwood Modifier
Forest/Woodland	East Gulf Coastal Plain Northern Mesic Hardwood Forest
Forest/Woodland	East Gulf Coastal Plain Southern Loess Bluff Forest
Forest/Woodland	East Gulf Coastal Plain Southern Mesic Slope Forest
Forest/Woodland	Northeastern Interior Dry Oak Forest - Mixed Modifier
Forest/Woodland	Northeastern Interior Dry Oak Forest-Hardwood Modifier
Forest/Woodland	Northern Atlantic Coastal Plain Dry Hardwood Forest
Forest/Woodland	Ridge and Valley Calcareous Valley Bottom Glade and Woodland
Forest/Woodland	South-Central Interior Mesophytic Forest
Forest/Woodland	Southern and Central Appalachian Cove Forest
Forest/Woodland	Southern and Central Appalachian Oak Forest
Forest/Woodland	Southern and Central Appalachian Oak Forest - Xeric

Forest/Woodland	Southern Appalachian Montane Pine Forest and Woodland
Forest/Woodland	Southern Coastal Plain Dry Upland Hardwood Forest
Forest/Woodland	Southern Interior Low Plateau Dry-Mesic Oak Forest
Forest/Woodland	Southern Interior Low Plateau Dry-Mesic Oak Forest - Evergreen Modifier
Forest/Woodland	Southern Piedmont Dry Oak-(Pine) Forest - Hardwood Modifier
Forest/Woodland	Southern Piedmont Dry Oak-(Pine) Forest - Mixed Modifier
Forest/Woodland	Southern Piedmont Dry Oak-Heath Forest - Hardwood Modifier
Forest/Woodland	Southern Piedmont Dry Oak-Heath Forest - Mixed Modifier
Forest/Woodland	Southern Piedmont Dry Oak-Heath Forest - Virginia/Pitch Pine Modifier
Forest/Woodland	Southern Piedmont Mesic Forest
Forest/Woodland	Southern Ridge and Valley Dry Calcareous Forest
Forest/Woodland	Southern Ridge and Valley Dry Calcareous Forest - Hardwood Modifier
Wetlands	Atlantic Coastal Plain Blackwater Stream Floodplain Forest - Forest Modifier
Wetlands	Atlantic Coastal Plain Brownwater Stream Floodplain Forest
Wetlands	Atlantic Coastal Plain Nonriverine Swamp and Wet Hardwood Forest - Taxodium/Nyssa Modifier
Wetlands	Atlantic Coastal Plain Nonriverine Swamp and Wet Hardwood Forest - Oak Dominated Modifier
Wetlands	Atlantic Coastal Plain Northern Basin Peat Swamp
Wetlands	Atlantic Coastal Plain Small Blackwater River Floodplain Forest
Wetlands	Atlantic Coastal Plain Small Brownwater River Floodplain Forest
Wetlands	Central Appalachian Floodplain - Forest Modifier
Wetlands	Central Appalachian Riparian - Forest Modifier
Wetlands	East Gulf Coastal Plain Large River Floodplain Forest - Forest Modifier
Wetlands	East Gulf Coastal Plain Small Stream and River Floodplain Forest
Wetlands	Lower Mississippi River Bottomland and Floodplain Forest
Wetlands	Lower Mississippi River Bottomland Depressions - Forest Modifier
Wetlands	Mississippi River Low Floodplain (Bottomland) Forest
Wetlands	Mississippi River Riparian Forest
Wetlands	North-Central Appalachian Acidic Swamp
Wetlands	North-Central Interior and Appalachian Rich Swamp
Wetlands	South-Central Interior Large Floodplain - Forest Modifier
Wetlands	South-Central Interior Small Stream and Riparian
Wetlands	Southern Coastal Plain Blackwater River Floodplain Forest
Wetlands	Southern Piedmont Large Floodplain Forest - Forest Modifier
Wetlands	Southern Piedmont Small Floodplain and Riparian Forest

**CITATIONS:** American Ornithologists' Union (AOU), Committee on Classification and Nomenclature. 1983. Check-list of North American Birds. Sixth Edition. American Ornithologists' Union, Allen Press, Inc., Lawrence, Kansas.

Barlow, J. C. 1980. Patterns of ecological interactions among migrant and resident vireos on the wintering grounds. Pages 79-107 in B80KEA02NA.

Bent, A.C. 1950. Life histories of North American wagtails, shrikes, vireos, and their allies. U.S. Natl. Mus. Bull. 197. Washington, D.C.

Bushman, E.S., and G.D. Therres. 1988. Habitat management guidelines for forest interior breeding birds of coastal Maryland. Maryland Department of Natural Resources, Wildlife Tech. Publ. 88-1. 50 pp.

DeGraaf, R.M., and J.H. Rappole. 1995. Neotropical migratory birds:natural history, distribution, and population change. Comstock Publishing Associates, Ithaca, NY.

Fussell, J. III and M. Lyons. 1990. Birds of the Outer Banks [pamphlet]. Eastern National Parks and Monument Association Coastal Wildlife Refuge Society.

Fussell, J.O. III. 1994. A birder's guide to coastal North Carolina. Chapel Hill and London: The University of North Carolina Press.

Hamel, P. B. 1992. The land manager's guide to the birds of the south. The Nature Conservancy, Chapel Hill, North Carolina. 367 pp + several appendices.

Hammerson, G. 2004. Yellow-throated Vireo (*Vireo flavifrons*), Ecology & Life History,Habitat Comments, in NatureServe Explorer website: <http://www.natureserve.org/explorer/>

Harrison, C. 1978. A field guide to the nests, eggs and nestlings of North American birds. Collins, Cleveland, Ohio.

Harrison, H.H. 1975. A field guide to bird's nests in the U.S. east of the Mississippi River. Houghton Mifflin Company, Boston, Massachusetts. 257 p.

- Harrison, H.H. 1979. A field guide to western birds' nests. Houghton Mifflin Company, Boston. 279 pp.
- Hilty, S.L., and W.L. Brown. 1986. A guide to the birds of Colombia. Princeton University Press, Princeton, New Jersey. 836 pp.
- James, R. D. 1973. Ethological and ecological relationships of yellow-throated and solitary vireos (Aves: Vireonidae) in Ontario. Ph.D. thesis, Department of Zoology, University of Toronto, Toronto, Ontario, Canada.
- Johnson, N. K., R. M. Zink, and J. A. Marten. 1988. Genetic evidence for relationships in the avian family Vireonidae. *Condor* 90:428-445.
- Kaufman K. 1996. Lives of North American Birds. Boston, New York: Houghton Mifflin Company.
- Keast, A., and E. S. Morton. 1980. Migrant birds in the Neotropics; ecology, distribution, and conservation. Smithsonian Inst. Press, Washington, D.C.
- McKenzie, P. M., and R. E. Noble. 1989. Sight records for Connecticut warbler (*OPORORNIS AGILIS*) and yellow-throated vireo (*VIREO FLAVIFRONS*) in Puerto Rico. *Florida Field Nat.* 17:69-72.
- Murray, B. W., et al. 1994. The use of cytochrome B sequence variation in estimation of phylogeny in the Vireonidae. *Condor* 96:1037-1054.
- Nicholson CP. 1997. Atlas of the breeding birds of Tennessee. Knoxville: University of Tennessee Press.
- Palmer-Ball, B.L., Jr. 1996. The Kentucky Breeding Bird Atlas. The University Press of Kentucky, Lexington.
- Potter, E. F., J. F. Parnell, and R. P. Teulings. 1980. Birds of the Carolinas. Univ. North Carolina Press, Chapel Hill. 408 pp.
- Ridgely, R.S., and G. Tudor. 1989. The birds of South America. Vol. 1. The Oscine passerines. Univ. Texas Press, Austin. 516 pp.
- Rodewald, P.G. and R.D. James. 1996. Yellow-throated vireo (*Vireo flavifrons*). In A. Poole and F. Gill, eds., *The Birds of North America*, No. 247. The Academy of Natural Sciences, Philadelphia and The American Ornithologists' Union, Washington, DC.
- Simpson MB Jr. 1992. Birds of the Blue Ridge Mountains. Chapel Hill and London: University of North Carolina Press.
- Stiles, F.G., and A.F. Skutch. 1989. A guide to the birds of Costa Rica. Comstock Publ. Associates, Cornell University Press, Ithaca, New York. 511 pp.
- Terres, J.K. 1980. The Audubon Society encyclopedia of North American birds. Alfred A. Knopf, New York.
- Whitcomb, R. F., C. S. Robbins, J. F. Lynch, B. L. Whitcomb, M. K. Klimciewicz, and D. Bystrak. 1981. Effects of forest fragmentation on avifauna of the eastern deciduous forest. Pages 125-206 in R. L. Burgess, and B. L. Sharpe (editors). *Forest island dy*

---

For more information:: SE-GAP Analysis Project / BaSIC  
 127 David Clark Labs  
 Dept. of Biology, NCSU  
 Raleigh, NC 27695-7617  
 (919) 513-2853  
[www.basic.ncsu.edu/segap](http://www.basic.ncsu.edu/segap)

Compiled: 15 September 2011

This data was compiled and/or developed by the Southeast GAP Analysis Project at The Biodiversity and Spatial Information Center, North Carolina State University.